


**(METRICALLY) QUARTER-STRATIFIABLE
SPACES AND THEIR APPLICATIONS
IN THE THEORY OF SEPARATELY CONTINUOUS FUNCTIONS**

TARAS BANAKH

ABSTRACT. We introduce and study (metrically) quarter-stratifiable spaces and then apply them to generalize Rudin and Kuratowski-Montgomery theorems about the Baire and Borel complexity of separately continuous functions.

The starting point for writing this paper was the desire to improve the results of V.K. Maslyuchenko et al. [MMMS], [MS], [KM], [KMM] who generalized a classical theorem of W.Rudin [Ru] which states that every separately continuous function $f : X \times Y \rightarrow \mathbb{R}$ on the product of a metrizable space X and a topological space Y belongs to the first Baire class. It was proven in [MMMS] that the metrizability of X in the Rudin theorem can be weakened to the σ -metrizability and paracompactness of X . A subtle analysis of Rudin's original proof reveals that this theorem is still valid for a much wider class of spaces X . These spaces are of independent interest, so we decided to give them a special name — metrically quarter-stratifiable spaces. (Metrically) quarter-stratifiable spaces are introduced and studied in details in the first three sections of this paper, where we investigate relationships between the class of (metrically) quarter-stratifiable spaces and other known classes of generalized metric spaces. It turns out that each semi-stratifiable space is quarter-stratifiable (this is a reason for the choice of the term “quarter-stratifiable”), while each quarter-stratifiable Hausdorff space has G_δ -diagonal. Because of this, the class of quarter-stratifiable spaces is “orthogonal” to the class of compacta — their intersection contains only metrizable compacta. The class of quarter-stratifiable spaces is quite wide and has many nice inheritance properties. Moreover, every (submetrizable) space with G_δ -diagonal is homeomorphic to a closed subset of a (metrically) quarter-stratifiable T_1 -space. The following diagram describes the interplay between the class of (metrically) quarter-stratifiable spaces and other classes of generalized metric spaces in the framework of Hausdorff spaces.



In the last three sections we apply metrically quarter-stratifiable spaces for generalizing classical theorems of Rudin [Ru] and Kuratowski-Montgomery [Ku₁], [Ku₂], [Mo].

1. QUARTER-STRATIFIABLE SPACES

Before introducing quarter-stratifiable spaces we recall some concepts from the theory of generalized metric spaces, see [Gr]. All topological spaces considered in this paper are T_1 -spaces, all maps (unlike to functions) are continuous.

1.1. Definition. A topological space X is

- (1) *perfect* if each open set in X is an F_σ -set;
- (2) *submetrizable* if X admits a *condensation* (i.e., a bijective continuous map) onto a metrizable space;
- (3) *σ -metrizable* if X is covered by a countable collection of closed metrizable subspaces;
- (4) a *G_δ -diagonal* if the diagonal is a G_δ -set in the square $X \times X$;
- (5) a *σ -space* if X has a σ -discrete network;
- (6) *developable* if there exists a sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of open covers of X such that for each $x \in X$ the set $\{\mathcal{St}(x, \mathcal{U}_n)\}_{n \in \mathbb{N}}$ is a base at x , where $\mathcal{St}(x, \mathcal{U}_n) = \cup\{U \in \mathcal{U}_n : x \in U\}$;
- (7) a *Moore space* if X is regular and developable;
- (8) *semi-stratifiable* (resp. *stratifiable*) if there exists a function G which assigns to each $n \in \mathbb{N}$ and a closed subset $H \subset X$, an open set $G(n, H)$ containing H such that
 - (i) $H = \bigcap_{n \in \mathbb{N}} G(n, U)$ (resp. $H = \bigcap_{n \in \mathbb{N}} \overline{G(n, U)}$) and
 - (ii) $G(n, K) \supset G(n, H)$ for every closed subset $K \supset H$ and $n \in \mathbb{N}$.

Semi-stratifiable spaces admit the following characterization, see [Gr, 5.8].

1.2. Theorem. A topological space (X, τ) is semi-stratifiable if and only if there exists a function $g : \mathbb{N} \times X \rightarrow \tau$ such that for every $x \in X$

- (1) $\{x\} = \bigcap_{n \in \mathbb{N}} g(n, x)$
- (2) $(x \in g(n, x_n), n \in \mathbb{N}) \Rightarrow (x_n \rightarrow x)$.

Weakening the first condition to $\bigcup_{x \in X} g(n, x) = X$ leads to the definition of a quarter-stratifiable space — the principal concept in this paper.

1.3. Definition. A topological space (X, τ) is called *quarter-stratifiable* if there exists a function $g : \mathbb{N} \times X \rightarrow \tau$ (called *quarter-stratifying function*) such that

- (1) $X = \bigcup_{x \in X} g(n, x)$ for every $n \in \mathbb{N}$;
- (2) $(x \in g(n, x_n), n \in \mathbb{N}) \Rightarrow (x_n \rightarrow x)$.

A space X is *hereditarily quarter-stratifiable* if every subspace of X is quarter-stratifiable.

Since the semi-stratifiability is a hereditary property, we get that each semi-stratifiable space is hereditarily quarter-stratifiable. As we shall see later, the converse is not true: the Sorgenfrey arrow, being hereditarily quarter-stratifiable, is not semi-stratifiable.

First, we give characterizations of the quarter-stratifiability in terms of open covers as well as of lower semi-continuous multivalued functions. We recall that a multivalued function $\mathcal{F} : X \rightarrow Y$ is *lower semi-continuous* if for every open set $U \subset Y$ its “preimage” $\mathcal{F}^{-1}(U) = \{x \in X : \mathcal{F}(x) \cap U \neq \emptyset\}$ is open in X . For a topological space X by X_d we denote X endowed with the discrete topology.

1.4. Theorem. *For a space X the following statements are equivalent:*

- (1) X is quarter-stratifiable;
- (2) there exists a sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of open covers of X and a sequence $\{s_n : \mathcal{U}_n \rightarrow X\}_{n \in \mathbb{N}}$ of functions such that $s_n(U_n) \rightarrow x$ if $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$;
- (3) there exists a sequence $\{\mathcal{F}_n : X \rightarrow X_d\}_{n \in \mathbb{N}}$ of lower semi-continuous multivalued functions tending to the identity map of X in the sense that for every point $x \in X$ and every neighborhood $O(x) \subset X$ of x there is $n_0 \in \mathbb{N}$ such that $\mathcal{F}_n(x) \subset O(x)$ for every $n \geq n_0$.

Proof. (1) $\Rightarrow$ (2) Assume (X, τ) is quarter-stratifiable and fix a function $g : \mathbb{N} \times X \rightarrow \tau$ from Definition 1.3. Let $\mathcal{U}_n = \{g(n, x) : x \in X\}$ and for every $U \in \mathcal{U}_n$ find $x \in X$ with $U = g(n, x)$ and let $s_n(U) = x$. Clearly that the so-defined sequences $\{\mathcal{U}_n\}$ and $\{s_n\}$ satisfy our requirements.

(2) $\Rightarrow$ (3) For every $n \in \mathbb{N}$ define $\mathcal{F}_n : X \rightarrow X_d$ letting $\mathcal{F}_n(x) = \{s_n(U) : x \in U \in \mathcal{U}_n\}$ for $x \in X$. Clearly, $\mathcal{F}_n$ is lower-semicontinuous and $\mathcal{F}_n(x) \rightarrow x$ for each $x \in X$.

(3) $\Rightarrow$ (1) For each $(n, x) \in \mathbb{N} \times X$ let $g(n, x) = \mathcal{F}_n^{-1}(x)$ and check that the so-defined function $g : \mathbb{N} \times X \rightarrow \tau$ fits. $\square$

Given a topological space X by $l(X)$ and $d(X)$ we denote respectively the Lindelöf number and the density of X .

Like other classes of generalized metric spaces, the class of quarter-stratifiable spaces has many nice inheritance properties.

1.5. Theorem. *Let X be a quarter-stratifiable space.*

- (1) Every open subspace and every retract of X is quarter-stratifiable.
- (2) If X is Hausdorff, then it has a G_δ -diagonal and countable pseudo-character.
- (3) If X is Hausdorff, then every paracompact Čech-complete subspace and every countably compact subspace of X is metrizable.

- (4) $d(X) \leq l(X)$.
- (5) If $f : Y \rightarrow X$ is a condensation with sequentially continuous inverse f^{-1} , then Y is quarter-stratifiable.
- (6) If $f : X \rightarrow Y$ is a finite-to-one open surjective map, then the space Y is quarter-stratifiable.

Proof. Using Theorem 1.4, fix a map $g : \mathbb{N} \times X \rightarrow \tau$, a sequence $\{\mathcal{F}_n : X \rightarrow X_d\}_{n \in \mathbb{N}}$ of lower-semicontinuous multivalued functions tending to the identity map of X , and sequences $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of open covers and functions $\{s_n\}_{n \in \mathbb{N}}$ satisfying the conditions of Definition 1.3 and Theorem 1.4.

1) Given an open subspace Y of X , let $\mathcal{V}_n = \{U \cap Y : U \in \mathcal{U}_n\}$, $n \in \mathbb{N}$. For each $V \in \mathcal{V}_n$ find a set $U(V) \in \mathcal{U}_n$ with $V = U(V) \cap Y$ and let $\tilde{s}_n(V) = s_n(U(V))$ if $s_n(U) \in Y$ and $\tilde{s}_n(V)$ be any point of Y , otherwise. Clearly, $\mathcal{V}_n$ is an open cover of Y . Now fix any point y of Y and take a sequence $V_n \in \mathcal{V}_n$ with $y \in V_n$ for $n \in \mathbb{N}$. Since $s_n(U(V_n)) \rightarrow y$, we get $s_n(U(V_n)) \in Y$ for all n beginning from some n_0 . Then $\tilde{s}_n(V_n) = s_n(U(V_n))$ for $n \geq n_0$ and thus $\tilde{s}_n(V_n) \rightarrow y$, which proves that Y is a quarter-stratifiable space.

Now suppose that a subspace Y of X is a retract of X and let $r : X \rightarrow Y$ be the corresponding retraction. One can readily prove that $\{r \circ \mathcal{F}_n|_Y : Y \rightarrow Y_d\}$ is a sequence of lower-semicontinuous multivalued functions tending to the identity map of Y .

2) Assume that the quarter-stratifiable space X is Hausdorff. Let $G_n = \bigcup_{U \in \mathcal{U}_n} U \times U \subset X \times X$. We claim that $\bigcap_{n \in \mathbb{N}} G_n$ coincides with the diagonal of $X \times X$. To show this, take any distinct points $x, y \in X$ and take disjoint neighborhood $O(x), O(y)$ of x and y , respectively. It follows from the choice of the sequences $\{\mathcal{U}_n\}$ and $\{s_n\}$ that there is $n \in \mathbb{N}$ such that $s_m(U_m) \in O(x)$ and $s_m(V_m) \in O(y)$ for all $m \geq n$ and $x \in U_m \in \mathcal{U}_m$, $y \in V_m \in \mathcal{U}_m$. We claim that $(x, y) \notin G_n$. Assuming the converse, we would find a set $U_n \in \mathcal{U}_n$ with $(x, y) \in U_n \times U_n$. Then $x, y \in U_n$ and $s_n(U_n) \in O(x) \cap O(y)$, a contradiction. Thus X has a G_δ -diagonal, which implies the countability of the pseudo-character of X .

3) If X is Hausdorff and $Y \subset X$ is a paracompact Čech-complete or countably compact subspace of X , then Y has a G_δ -diagonal and by [Bo, 8.2] or [Gr, 2.14] is metrizable.

4) To see that $d(X) \leq l(X)$ we may take the subset $D = \{s_n(U) : U \in \mathcal{V}_n, n \in \mathbb{N}\}$, where $\mathcal{V}_n$ is a subcover of $\mathcal{U}_n$ of size $|\mathcal{V}_n| \leq l(X)$ for $n \in \mathbb{N}$. Clearly, D is a dense subset in X of size $|D| \leq l(X)$.

5) If $f : Y \rightarrow X$ is a condensation with sequentially continuous inverse f^{-1} , we may consider the function g_Y assigning to each pair $(n, y) \in \mathbb{N} \times Y$ the open set $f^{-1}(g(n, f(y)))$ of Y . One may easily show that the so-defined function g_Y turns Y into a quarter-stratifiable space.

6) Suppose $f : X \rightarrow Y$ is an open finite-to-one surjective map. It can be shown that for every $n \in \mathbb{N}$ the multivalued map $f \circ \mathcal{F}_n \circ f^{-1} : Y \rightarrow Y_d$ defined by $f \circ \mathcal{F}_n \circ f^{-1} = \bigcup_{x \in f^{-1}(y)} f(\mathcal{F}_n(x))$ for $y \in Y$ is lower-semicontinuous. Moreover, the sequence $\{f \circ \mathcal{F}_n \circ f^{-1}\}_{n \in \mathbb{N}}$ tends to the identity map of Y . Then according to Theorem 1.4, the space Y is quarter-stratifiable. $\square$

Next, we show that the quarter-stratifiability is stable with respect to certain set-theoretic operations.

1.6. Theorem.

- (1) *The countable product of quarter-stratifiable spaces is quarter-stratifiable.*
- (2) *A space X is quarter-stratifiable, provided $X = A \cup B$ is a union of two quarter-stratifiable subspaces of X , one of which is a closed G_δ -set in X .*
- (3) *A space X is quarter-stratifiable, provided X can be covered by a σ -locally finite collection of quarter-stratifiable closed G_δ -subspaces of X .*

Proof.

1) Let X_i , $i \in \mathbb{N}$, be a sequence of quarter-stratifiable spaces. For every $i \in \mathbb{N}$, fix a point $*_i \in X_i$ and a sequence $\{\mathcal{F}_{i,n} : X_i \rightarrow (X_i)_d\}_{n \in \mathbb{N}}$ of lower-semicontinuous multivalued functions tending to the identity map of X_i as $n \rightarrow \infty$. Let $X = \prod_{i \in \mathbb{N}} X_i$ and for every $n \in \mathbb{N}$ identify the product $\prod_{i=1}^n X_i$ with the subspace $\{(x_i)_{i \in \mathbb{N}} : x_i = *_i \text{ for } i > n\}$ of X . For every $n \in \mathbb{N}$ define a multivalued function $\mathcal{F}_n : X \rightarrow X_d$ as follows: $\mathcal{F}_n((x_i)_{i \in \mathbb{N}}) = \prod_{i=1}^n \mathcal{F}_{i,n}(x_i)$. It is easy to see that each function $\mathcal{F}_n$ is lower semicontinuous and the sequence $\{\mathcal{F}_n\}$ tends to id_X .

2) Suppose $X = A \cup B$, where A, B are quarter-stratifiable subspaces and B is a closed G_δ -set in X . Fix a decreasing sequence $\{O_n(B)\}_{n \in \mathbb{N}}$ of open subsets of X with $B = \bigcap_{n \in \mathbb{N}} O_n(B)$. By Theorem 1.5(1), the open subspace $A \setminus B = X \setminus B$ of A is quarter-stratifiable and thus admits a function $g_{A \setminus B} : \mathbb{N} \times (A \setminus B) \rightarrow \tau$ into the topology τ of X such that $\bigcup_{x \in A \setminus B} g_{A \setminus B}(n, x) \supset A \setminus B$ for each $n \in \mathbb{N}$, and $x \in g_{A \setminus B}(n, x_n) \Rightarrow x_n \rightarrow x$ for each $x, x_n \in A \setminus B$, $n \in \mathbb{N}$.

Using the quarter-stratifiability of B find a function $g_B : \mathbb{N} \times B \rightarrow \tau$ such that $B \subset \bigcup_{x \in B} g_B(n, x) \subset O_n(B)$ for every $n \in \mathbb{N}$, and $x \in g_B(n, x_n) \Rightarrow x_n \rightarrow x$ for each $x, x_n \in B$, $n \in \mathbb{N}$. It is easy to verify that the function $g : \mathbb{N} \times X \rightarrow \tau$ defined by

$$g(n, x) = \begin{cases} g_{A \setminus B}(n, x) & \text{if } x \in A \setminus B, \\ g_B(n, x) & \text{if } x \in B \end{cases}$$

turns X into a quarter-stratifiable space.

3) Let $\{X_i\}_{i \in \mathcal{I}}$ be a σ -locally finite collection of quarter-stratifiable closed G_δ -subspaces of a space X . Write $\mathcal{I} = \bigcup_{k \in \mathbb{N}} \mathcal{I}_k$ so that the collection $\{X_i\}_{i \in \mathcal{I}_k}$ is locally finite for every $k \in \mathbb{N}$. Without loss of generality, $\mathcal{I}_k \subset \mathcal{I}_{k+1}$ for every k . For every k fix an open cover $\mathcal{W}_k$ of X whose any element $W \in \mathcal{W}_k$ meets only finitely many of the sets X_i 's with $i \in \mathcal{I}_k$. We may additionally require that the cover $\mathcal{W}_k$ is inscribed into the cover $\mathcal{W}_{k-1}$ for $k > 1$.

Since X_i 's are G_δ -sets in X , for every $i \in \mathcal{I}$ we may find a decreasing sequence $\{O_n(X_i)\}_{n \in \mathbb{N}}$ of open subsets $O_n(X_i) \subset \mathcal{St}(X_i, \mathcal{W}_n) = \{W \in \mathcal{W}_n : W \cap X_i \neq \emptyset\}$ with $\bigcap_{n \in \mathbb{N}} O_n(X_i) = X_i$.

Let $\leq$ be any well-ordering of the index set $\mathcal{I}$ such that $i < j$ for every $i, j \in \mathcal{I}$ with $i \in \mathcal{I}_k \not\subset \mathcal{I}_j$ for some k . It follows from the local finiteness of the collections $\{X_i\}_{i \in \mathcal{I}_k}$ that for every $i \in \mathcal{I}$ the set $\bigcup_{j < i} X_j$ is closed in X . Using the quarter-stratifiability of open subsets of X_i 's, for every $i \in \mathbb{N}$ fix a sequence $\{\mathcal{U}_{i,n}\}_{n \in \mathbb{N}}$ of open covers of the set $Y_i = X_i \setminus \bigcup_{j < i} X_j$ and a sequence $\{s_{i,n} : \mathcal{U}_{i,n} \rightarrow Y_i\}$ of functions such that $s_{i,n}(U_n) \rightarrow y$ for every $Y_i \ni y \in U_n \in \mathcal{U}_{i,n}$, $n \in \mathbb{N}$. Without loss of generality, we may assume that each $U \in \mathcal{U}_{i,n}$ is an open subset of X lying in $O_n(X_i) \setminus \bigcup_{j < i} X_j$. Now for every $n \in \mathbb{N}$ consider the open cover $\mathcal{U}_n = \bigcup_{i \in \mathcal{I}} \mathcal{U}_{i,n}$ and the function $s_n = \bigcup_{i \in \mathcal{I}} s_{i,n} : \mathcal{U}_n \rightarrow X$.

To show that the space X is quarter-stratifiable, it suffices to verify that for every $x \in X$ and a sequence $U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$, with $x \in U_n$ we have $s_n(U_n) \rightarrow x$. Let $i = \min\{j \in \mathcal{I} : x \in X_j\}$. Then $x \in Y_i$. For every $n \in \mathbb{N}$ find $i_n \in \mathcal{I}$ with $U_n \in \mathcal{U}_{i_n, n}$. Since $x \in U_n \subset O_{i_n}(X_{i_n}) \setminus \bigcup_{j < i_n} X_j$, we get $i_n \leq i$ for all $n \in \mathbb{N}$. It follows from the choice of the cover $\mathcal{W}_n$ and the sets $O_n(X_j)$ that $\bigcap_{n=1}^{\infty} (\bigcup_{j < i} O_n(X_j)) = \bigcup_{j < i} X_j$. Consequently, $x \notin \bigcap_{n=1}^{\infty} (\bigcup_{j < i} O_n(X_j))$ and there is $n_0 \in \mathbb{N}$ such that $x \notin \bigcup_{j < i} O_n(X_j)$ for every $n \geq n_0$. Since $x \in U_n \subset O_n(X_{i_n})$ for every $n \in \mathbb{N}$, we conclude that $i_n = i$ for $n \geq n_0$. Then $x \in U_n \in \mathcal{U}_{i, n}$ for all $n \geq n_0$ and $s_n(U_n) = s_{i, n}(U_n) \rightarrow x$, which completes the proof of the quarter-stratifiability of X . $\square$

1.7. Remark. The G_δ -condition in Theorem 1.6(2) is essential: the one-point compactification $\alpha\Gamma$ of an uncountable discrete space Γ may be written as the union $\alpha\Gamma = \{\infty\} \cup \Gamma$ of two metrizable subspaces, one of which is closed. But $\alpha\Gamma$, being a non-metrizable compactum, is not quarter-stratifiable, see Theorem 1.5(3). Yet, we do not know the answer to the following question.

1.8. Question. *Is a (regular) space X quarter-stratifiable if it is a union of two closed quarter-stratifiable subspaces?*

According to Theorem 1.5(1), every open subspace of a quarter-stratifiable space is quarter-stratifiable. It is not true for closed subspaces of quarter-stratifiable spaces: every G_δ -diagonal space is homeomorphic to a closed subset of a quarter-stratifiable T_1 -space.

1.9. Theorem. *Every space X with G_δ -diagonal is homeomorphic to a closed subspace of a quarter-stratifiable T_1 -space Y with $|Y \setminus X| \leq \max\{l(X), d(X)\}$.*

Proof. Suppose X is a space with G_δ -diagonal. By Theorem 2.2 of [Gr], X admits a sequence $\{\mathcal{V}_n\}_{n \in \mathbb{N}}$ of covers such that $\{x\} = \bigcap_{n \in \mathbb{N}} \text{St}(x, \mathcal{V}_n)$ for every $x \in X$. Replacing each $\mathcal{V}_n$ by a suitable subcover, we may assume that $|\mathcal{V}_n| \leq l(X)$ for every n , and each cover $\mathcal{V}_n$ consists of non-empty subsets of X . Let $D \subset X$ be a dense subset of X with $|D| \leq d(X)$. For every $n \in \mathbb{N}$ and every $V \in \mathcal{V}_n$ pick a point $c(V) \in V \cap D$. Let $S_n = \{0\} \cup \{1/i : i \geq n\}$ denote a tail of the convergent sequence S_1 .

Consider the subset $Y = X \times \{0\} \cup \{(c(V), 1/n) : V \in \mathcal{V}_n, n \in \mathbb{N}\} \subset X \times S_1$. It is clear that $|Y \setminus X| \leq \min\{|\bigcup_{n \in \mathbb{N}} \mathcal{V}_n|, |D|\} \leq \min\{l(X), d(X)\}$. Identify X with the subset $X \times \{0\}$ of Y . Define a topology τ on Y letting $U \subset Y$ be open if and only if $U \cap X$ is open in X and for every $x \in U \cap X$ there is $n_0 \in \mathbb{N}$ such that $\{(c(V), 1/n) : x \in V \in \mathcal{V}_n, n \geq n_0\} \subset U$. Thus X is homeomorphic to a closed subspace of Y , while all points of $Y \setminus X$ are isolated. Since each one-point subset of Y is closed, Y is a T_1 -space. To show that Y is a quarter-stratifiable space, consider open covers $\mathcal{U}_n = \{\{y\}, (V \times S_n) \cap Y : y \in Y \setminus X, V \in \mathcal{V}_n\}$, $n \in \mathbb{N}$, of Y . For every $U \in \mathcal{U}_n$ let

$$s_n(U) = \begin{cases} y & \text{if } U = \{y\} \text{ for some } y \in Y \setminus X; \\ (c(V), 1/n) & \text{if } U = (V \times S_n) \cap Y \text{ for } V \in \mathcal{V}_n. \end{cases}$$

It follows from the choice of the topology τ on Y that $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$, implies $s_n(U_n) \rightarrow x$ i.e., Y is a quarter-stratifiable T_1 -space. $\square$

1.10. Problem. *Describe the class of subspaces of regular (Tychonoff) quarter-stratifiable T_1 -spaces.*

2. METRICALLY QUARTER-STRATIFIABLE SPACES

In this section we introduce and study metrically quarter-stratifiable spaces forming a class, intermediate between the class of paracompact quarter-stratifiable Hausdorff spaces and the class of submetrizable quarter-stratifiable spaces. We start with defining a quarter-stratifying topology.

2.1. Definition. A topology τ' on a topological space (X, τ) is called *quarter-stratifying* if there exists a quarter-stratifying function $g : \mathbb{N} \times X \rightarrow \tau$ for X such that $g(\mathbb{N} \times X) \subset \tau'$.

A topological space X is defined to be *metrically quarter-stratifiable* if it admits a weaker metrizable quarter-stratifying topology.

The following theorem characterizes metrically quarter-stratifiable spaces.

2.2. Theorem. *For a space X the following statements are equivalent:*

- (1) X admits a weaker metrizable quarter-stratifying topology;
- (2) X admits a weaker paracompact Hausdorff quarter-stratifying topology;
- (3) there exists a weaker metrizable topology τ_m on X , a sequence $\{U_n\}_{n \in \mathbb{N}}$ of covers of X by τ_m -open subsets and a sequence $\{s_n : \mathcal{U}_n \rightarrow X\}_{n \in \mathbb{N}}$ of functions such that $s_n(U_n) \rightarrow x$ in X if $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$;
- (4) there exists a weaker paracompact Hausdorff topology τ_p on X , a sequence $\{V_n\}_{n \in \mathbb{N}}$ of covers of X by τ_p -open subsets and a sequence $\{\tilde{s}_n : \mathcal{V}_n \rightarrow X\}_{n \in \mathbb{N}}$ of functions such that $\tilde{s}_n(V_n) \rightarrow x$ in X if $x \in V_n \in \mathcal{V}_n$, $n \in \mathbb{N}$;

Proof. The equivalence (1) $\Leftrightarrow$ (3) and (2) $\Leftrightarrow$ (4) can be proved by analogy with the corresponding equivalences in Theorem 1.4; the implication (3) $\Rightarrow$ (4) is trivial.

(4) $\Rightarrow$ (3) Assume that τ_p is a weaker paracompact Hausdorff topology on X , $\{U_n\}_{n \in \mathbb{N}}$ is a sequence of τ_p -open covers of X , and $\{s_n : \mathcal{U}_n \rightarrow X\}_{n \in \mathbb{N}}$ is a sequence of functions such that $s_n(U_n) \rightarrow x$ in X if $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$. Let $l_1(\mathcal{U}_n)$ denote the Banach space of all absolutely summable functions $f : \mathcal{U}_n \rightarrow \mathbb{R}$ equipped with the norm $\|f\| = \sum_{U \in \mathcal{U}_n} |f(U)|$.

Using the paracompactness of the topology $\tau_p \subset \tau$, for every $n \in \mathbb{N}$ find a partition of unity $\{\lambda_U : X \rightarrow [0, 1]\}_{U \in \mathcal{U}_n}$ such that $\lambda_U^{-1}(0, 1] \subset U$ for $U \in \mathcal{U}_n$. Observe that this partition of unity can be seen as a continuous map $\Lambda_n : X \rightarrow l_1(\mathcal{U}_n)$ acting as $\Lambda_n(x) = (\lambda_U(x))_{U \in \mathcal{U}_n}$ (the continuity of Λ_n follows from the local finiteness of the cover $\{\lambda_U^{-1}(0, 1] : U \in \mathcal{U}_n\}$).

Let $\Lambda : X \rightarrow \prod_{n \in \mathbb{N}} l_1(\mathcal{U}_n)$ denote the diagonal product of the maps Λ_n . By analogy with the proof of Theorem 1.5(2), show that the map Λ is injective. Then the weakest topology τ_m on X for which the map Λ is continuous is metrizable. Observe that for each $n \in \mathbb{N}$ and $U \in \mathcal{U}_n$ the set $V(U) = \lambda_U^{-1}(0, 1]$ is τ_m -open. Let $\mathcal{V}_n = \{V(U) : U \in \mathcal{U}_n\}$. For every $V \in \mathcal{V}_n$ find $U \in \mathcal{U}_n$ with $V = V(U)$ and let $\tilde{s}_n(V) = s_n(U)$. One can easily check that the metrizable topology τ_m on X and the sequences $\{V_n\}_{n \in \mathbb{N}}$, $\{\tilde{s}_n\}_{n \in \mathbb{N}}$ satisfy the condition (3). $\square$

In the following theorem we collect some elementary properties of metrically quarter-stratifiable spaces.

2.3. Theorem.

- (1) *Each paracompact Hausdorff quarter-stratifiable space is metrically quarter-stratifiable.*
- (2) *Each metrically quarter-stratifiable space is submetrizable and quarter-stratifiable.*
- (3) *Every open subspace of a metrically quarter-stratifiable space is metrically quarter-stratifiable.*

- (4) If X is a metrically quarter-stratifiable space and $f : Y \rightarrow X$ is a condensation with sequentially continuous inverse f^{-1} , then the space Y is metrically quarter-stratifiable.
- (5) The product $X = \prod_{n \in \mathbb{N}} X_n$ of a countable collection of metrically quarter-stratifiable spaces X_n , $n \in \mathbb{N}$, is a metrically quarter-stratifiable space.
- (6) A space X is metrically quarter-stratifiable if there exist a weaker paracompact Hausdorff topology τ_p and two subspaces $A, B \subset X$ such that $A \cup B = X$, A is a closed G_δ -set in (X, τ_p) and the topologies induced by the topology τ_p on A and B are quarter-stratifying.
- (7) A space X is metrically quarter-stratifiable if there exist a weaker paracompact Hausdorff topology τ_p on X and a σ -locally finite cover $\mathcal{C}$ of (X, τ_p) by closed G_δ -subspaces such that the topology induced by τ_p on each $C \in \mathcal{C}$ is quarter-stratifying (with respect to the original topology of C);
- (8) A space X is metrically quarter-stratifiable if there is a weaker metrizable topology τ_m on X and a cover $\mathcal{C}$ of X , well-ordered by the inclusion relation, such that the topology τ_m induces the original topology on each $C \in \mathcal{C}$;
- (9) Every submetrizable space is homeomorphic to a metrically quarter-stratifiable space.

Proof. All statements (except for (8)) easily follow from definitions or can be proved by analogy with the corresponding properties of quarter-stratifiable spaces.

To prove the eighth statement, fix a continuous metric d on X and a well-ordered (by the inclusion relation) cover $\mathcal{C}$ of X such that d induces the original topology on each $C \in \mathcal{C}$. Let $\mathcal{U}_n$ denotes the collection of all open $1/n$ -balls with respect to the metric d . For every $U \in \mathcal{U}_n$ let $C(U)$ be the smallest set $C \in \mathcal{C}$ meeting U and let $s_n(U) \in U \cap C(U)$ be any point.

Fix any $x \in X$ and a sequence $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$. We have to show that $s_n(U_n) \rightarrow x$. Let $C(x)$ be the smallest set $C \in \mathcal{C}$ containing the point x . Since $x \in C(x) \cap U_n$, we conclude that $s_n(U_n) \in C(x)$. By the choice of the sets U_n , $\text{diam}(U_n) \leq 2/n$. Then $d(x, s_n(U_n)) \leq \text{diam}(U_n) \leq 2/n$, $n \in \mathbb{N}$, and thus $s_n(U_n) \rightarrow x$ (because d induces the topology of $C(x)$). $\square$

According to Theorem 2.3(2) each metrically quarter-stratifiable space is submetrizable and quarter-stratifiable. We do not know if the converse is also true.

2.4. Question. *Is every submetrizable quarter-stratifiable space metrically quarter-stratifiable?*

3. SOME EXAMPLES

In this section we collect some examples exposing the difference between the class of (metrizable) quarter-stratifiable spaces and other classes of generalized metric spaces. It is known that every Moore space is semi-stratifiable and each collectively normal Moore space is metrizable, see [Gr]. Yet, there exist Moore spaces which are not submetrizable, see [Ny]. Thus we have

3.1. Example. *There exists a quarter-stratifiable (Moore) space which is not metrically quarter-stratifiable.*

Theorems 1.9 and 2.3(9) show that the class of quarter-stratifiable spaces is much wider than the class of semi-stratifiable spaces. Using these theorems many wild (metrically) quarter-stratifiable spaces may be constructed. But in general, so-constructed spaces are not hereditarily quarter-stratifiable.

3.2. Example. *The Sorgenfrey arrow Z is not semi-stratifiable but every subspace of Z is Lindelöf, separable, and metrically quarter-stratifiable.*

Proof. Recall that the Sorgenfrey arrow Z is the semi-interval $[0, 1)$ endowed with the topology generated by the base consisting of all semi-intervals $[a, b)$ where $0 \leq a < b \leq 1$, see [En₁, 1.2.2]. Since Z embeds into a linearly ordered space (“two arrows” of Aleksandrov [En₁, 3.10.C]), Z is monotonically normal, see [Gr, 5.21]. If Z would be semi-stratifiable, then Z , being monotonically normal, would be stratifiable according to Theorem 5.16 of [Gr]. Since each stratifiable space is a σ -space [Gr, 5.9] and Lindelöf σ -spaces have countable network [Gr, 4.4], this would imply that Z has countable network, which is a contradiction. Therefore, the Sorgenfrey arrow Z is not semi-stratifiable.

Next, we show that every subspace X of Z is quarter-stratifiable. For every $n \in \mathbb{N}$ consider the finite open cover $\mathcal{U}_n = \{X \cap [\frac{k-1}{n}, \frac{k}{n}) : 0 < k \leq n\}$ of X . For every element $U \in \mathcal{U}_n$ choose a point $s_n(U) \in X$ as follows. If the set $\uparrow U = \{x \in X : x \geq \sup U\}$ is not empty, then let $s_n(U)$ be any point of X with $s_n(U) < \inf \uparrow U + \frac{1}{n}$; otherwise, let $s_n(U)$ be any point of X . It can be shown that $s_n(U_n) \rightarrow x$ for every $x \in X$ and every sequence $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$. Thus the space X is quarter-stratifiable. It is well known that the Sorgenfrey arrow is hereditarily Lindelöf and hereditarily separable which implies that the space X is Lindelöf and separable. Then X is also paracompact ([En₁, 5.1.2]) and hence, being quarter-stratifiable, is metrically quarter-stratifiable. $\square$

Since the metrical quarter-stratifiability is productive, we get

3.3. Example. *The square of the Sorgenfrey arrow is metrically quarter-stratifiable but not normal.*

Another example of a metrically quarter-stratifiable non-normal space is the Nemytski plane. Unlike to (semi-)stratifiable spaces metrically quarter-stratifiable spaces need not be perfect (or paracompact).

3.4. Example. *A separable zero-dimensional metrically quarter-stratifiable Tychonov space which is neither perfect nor Lindelöf.*

Proof. Consider the Cantor cube $2^\omega = \{0, 1\}^\omega$, i.e., the set of all functions $\omega \rightarrow \{0, 1\}$. There is a natural partial order on 2^ω : $x \leq y$ iff $x(i) \leq y(i)$ for each $i \in \omega$. For $k \in \{0, 1\}$ let $2_k^\omega = \{(x_i)_{i \in \omega} : \exists n \in \omega \text{ with } x(i) = k \text{ for all } i \geq n\}$. For a point $x \in 2^\omega$ and $n \in \omega$ let $U(x, n) = \{y \in 2^\omega : y \geq x \text{ and } y(i) = x(i) \text{ for } i \leq n\}$. On the set 2^ω consider the topology τ generated by the base $\{U(x, n) : x \in 2^\omega, n \in \omega\}$. Clearly, the space $X = (2^\omega, \tau)$ is Tychonoff, zero-dimensional, and separable (2_1^ω is a countable dense set in X).

Let us show that the space X is metrically quarter-stratifiable. For every $n \in \omega$ identify 2^n with the subset $\{(x_i) \in 2^\omega : x_i = 0 \text{ for } i \geq n\} \subset 2^\omega$ and consider the finite cover $\mathcal{U}_n = \{U(x, n) : x \in 2^n\}$ of 2^ω . It is clear that each set $U(x, n) \in \mathcal{U}_n$ is open in the product topology of 2^ω which is metrizable. To each $U \in \mathcal{U}_n$ assign the point $s_n(U) = \max U \in 2_1^\omega$. One may easily verify that $s_n(U_n) \rightarrow x$ for every $x \in X$ and a sequence $x \in U_n \in \mathcal{U}_n$, $n \in \omega$. Thus the space X is metrically quarter-stratifiable.

Now we show that the space X is not perfect. Note that 2_0^ω is a closed subset of X . We claim that it is not a G_δ -subset of X . Assume conversely that $2_0^\omega = \bigcap_{n \in \mathbb{N}} O_n$, where O_n are open subsets of X . Since the topology of X coincides with the usual product topology at

every point of the set 2_0^ω , we may assume that each set O_n is open in the product topology of 2^ω . Then by the Baire theorem, the intersection $\bigcap_{n \in \mathbb{N}} O_n$ is a dense G_δ -subset in 2^ω , which contains a point $x \notin 2_0^\omega$, a contradiction.

By Theorem 2.3(3) the open subspace $Y = X \setminus 2_0^\omega$ of X is metrically quarter-stratifiable. Since X is regular and 2_0^ω is not a G_δ -set in X , we conclude that the space Y is not Lindelöf.

Then the product $X \times Y$ is a separable Tychonov zero-dimensional metrically quarter-stratifiable space which is neither perfect nor Lindelöf. $\square$

3.5. Question. *Is there a hereditarily quarter-stratifiable space which is not perfect?*

We recall that a topological space X is *finally compact* if $l(X) \leq \aleph_0$, i.e., every open cover of X admits a countable subcover. Finally compact regular spaces are called *Lindelöf*.

3.6. Example. *A hereditarily finally compact submetrizable uncountable space whose uncountable subspaces are neither separable nor quarter-stratifiable.*

Proof. Let X be the interval $(0, 1)$ with the topology generated by the sets of the form $(a, b) \setminus C$, where $0 \leq a < b \leq 1$ and C is a countable set. Clearly, the space X is submetrizable and hereditarily finally compact. Yet, every countable subset of X is closed which implies that every uncountable subspace Y of X is not separable. Since finally compact quarter-stratifiable spaces are separable, see Theorem 1.6(4), we conclude that the space Y is not quarter-stratifiable. $\square$

The space constructed in Example 3.6 is not regular. There is also a regular submetrizable non-quarter-stratifiable space.

Given a subset $B \subset \mathbb{R}$ denote by $\mathbb{R}_B$ the real line $\mathbb{R}$ endowed with the topology consisting of the sets $U \cup A$, where U open in $\mathbb{R}$ and $A \subset B$, see [En₁, 5.1.22 and 5.5.2]. It is known that for every $B \subset \mathbb{R}$ the space $\mathbb{R}_B$ is hereditarily paracompact; $\mathbb{R}_B$ is perfect if and only if B is a G_δ -set in $\mathbb{R}$. If $B \subset \mathbb{R}$ is a Bernstein set, then the space $\mathbb{R}_B$ is Lindelöf, see [En₁, 5.5.4]. Recall that a subset $B \subset \mathbb{R}$ is called a *Bernstein set* if $C \cap B \neq \emptyset \neq C \setminus B$ for every uncountable compactum $C \subset \mathbb{R}$. Bernstein sets can be easily constructed by transfinite induction, see [En₁, 5.4.4].

3.7. Example. *If $B \subset \mathbb{R}$ is not σ -compact, then the space $\mathbb{R}_B$ is not quarter-stratifiable. Consequently, if $B \subset \mathbb{R}$ is the set of irrationals, then $\mathbb{R}_B$ is a hereditarily paracompact submetrizable perfect zero-dimensional space which is neither separable nor quarter-stratifiable; if $B \subset \mathbb{R}$ is a Bernstein set, then $\mathbb{R}_B$ is a submetrizable hereditarily paracompact Lindelöf zero-dimensional space which is neither perfect nor separable nor quarter-stratifiable.*

Proof. Assume that $B \subset \mathbb{R}$ is not σ -compact. Suppose the space $\mathbb{R}_B$ is quarter-stratifiable and fix a sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of open covers of $\mathbb{R}_B$ and a sequence $\{s_n : \mathcal{U}_n \rightarrow \mathbb{R}_B\}_{n \in \mathbb{N}}$ of functions such that $s_n(U_n) \rightarrow x$ for every $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$. Since the real line is hereditarily Lindelöf and the topology of $\mathbb{R}_B$ coincides with the usual topology at points of the set $\mathbb{R} \setminus B$, we may find countable subcovers $\mathcal{V}_n \subset \mathcal{U}_n$, $n \in \mathbb{N}$, of the set $\mathbb{R} \setminus B$. Observe that $C = \{s_n(V) : V \in \mathcal{V}_n, n \in \mathbb{N}\}$ is countable while the intersection $G = \bigcap_{n \in \mathbb{N}} St(\mathbb{R} \setminus B, \mathcal{V}_n)$ is a G_δ -set in the usual topology of $\mathbb{R}$. Since B is not a σ -compact, there is a point $x \in B \cap G \setminus C$. Choose for every $n \in \mathbb{N}$ an element $V_n \in \mathcal{V}_n$ with $V_n \ni x$. Then $x \neq s_n(V_n)$ for all n and by the definition of the topology of $\mathbb{R}_B$, $s_n(V_n) \not\rightarrow x$. This contradiction shows that the space $\mathbb{R}_B$ is not quarter-stratifiable. $\square$

4. A GENERALIZATION OF KURATOWSKI-MONTGOMERY THEOREM

In this section we apply metrically quarter-stratifiable spaces to generalize a theorem of Kuratowski [Ku₂] and Montgomery [Mo] which asserts that for metrizable spaces X, Y, Z a function $f : X \times Y \rightarrow Z$ is Borel measurable of class $\alpha + 1$, α a countable ordinal, if f is continuous with respect to the first variable and Borel measurable of class α with respect to the second variable.

At first, we recall the definitions of the multiplicative and additive Borel classes. Given a topological space X let $\mathcal{A}_0(X)$ and $\mathcal{M}_0(X)$ denote the classes of all open and all closed subsets of X , respectively. Assuming that for a countable ordinal α the classes $\mathcal{A}_\beta(X)$, $\mathcal{M}_\beta(X)$, $\beta < \alpha$, are already defined, let $\mathcal{A}_\alpha(X) = \{\bigcup_{n=1}^\infty M_n : M_n \in \bigcup_{\beta < \alpha} \mathcal{M}_\beta(X) \text{ for all } n \in \mathbb{N}\}$ and $\mathcal{M}_\alpha(X) = \{M \subset X : X \setminus M \in \mathcal{A}_\alpha(X)\}$.

We say that a function $f : X \rightarrow Y$ between topological spaces is Borel measurable of class α if the preimage $f^{-1}(U)$ of any open set $U \subset Y$ belongs to the additive class $\mathcal{A}_\alpha(X)$ (of course, this is equivalent to saying that the preimage $f^{-1}(F)$ of any closed subset $F \subset Y$ belongs to the multiplicative class $\mathcal{M}_\alpha(X)$).

The set of all Borel measurable functions $f : X \rightarrow Y$ of class α is denoted by $H_\alpha(X, Y)$. For topological spaces X, Y, Z and a countable ordinal α let $CH_\alpha(X \times Y, Z)$ denote the set of all functions $f : X \times Y \rightarrow Z$ which are continuous with respect to the first variable and are Borel measurable of class α with respect to the second variable. The Kuratowski-Montgomery Theorem states that $CH_\alpha(X \times Y, Z) \subset H_{\alpha+1}(X \times Y, Z)$ for metrizable spaces X, Y, Z and a countable ordinal α .

A subset A of a topological space X is called a $\overline{G}_\delta$ -set if $A = \bigcap_{n \in \mathbb{N}} U_n = \bigcap_{n \in \mathbb{N}} \bar{U}_n$ for some open sets $U_n \subset X$, $n \in \mathbb{N}$. Observe that every $\overline{G}_\delta$ -set is a closed G_δ -set and every closed G_δ -set in a normal space is a $\overline{G}_\delta$ -set. Consequently, every closed subset of a perfectly normal space is a $\overline{G}_\delta$ -set.

4.1. Theorem. *The inclusion $CH_\alpha(X \times Y, Z) \subset H_{\alpha+1}(X \times Y, Z)$ holds for any countable ordinal α , any metrically quarter-stratifiable space X , any topological space Y , and any topological space Z whose every closed subset is a $\overline{G}_\delta$ -set.*

Proof. Using Theorem 2.2, fix a weaker metrizable topology τ_m on the metrically quarter-stratifiable space X , a sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of τ_m -open covers of X and a sequence $\{s_n : \mathcal{U}_n \rightarrow X\}_{n \in \mathbb{N}}$ of functions such that $s_n(U_n) \rightarrow x$ if $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$. According to the classical Stone Theorem [St], we may assume that each cover $\mathcal{U}_n$ is locally finite and σ -discrete.

Fix any function $f \in CH_\alpha(X \times Y, Z)$. To show that $f \in H_{\alpha+1}(X \times Y, Z)$ we have to verify that the preimage $f^{-1}(F)$ of any closed subset $F \subset Z$ belongs to the class $\mathcal{M}_{\alpha+1}(X \times Y)$.

The set F , being $\overline{G}_\delta$ -set, can be written as $F = \bigcap_{n=1}^\infty \overline{W}_n$, where $W_n \subset W_{n-1}$ are open neighborhoods of F . For every $n \in \mathbb{N}$ and $U \in \mathcal{U}_n$ let $f_U : Y \rightarrow Z$ be the map defined by $f_U(y) = f(s_n(U), y)$ for $y \in Y$. It is clear that $f_U \in H_\alpha(Y, Z)$ which yields $f_U^{-1}(W_m) \in \mathcal{A}_\alpha(Y)$ for every m . Since the set U is functionally open in X , we get $U \times f_U^{-1}(W_m) \in \mathcal{A}_\alpha(X \times Y)$, $m \in \mathbb{N}$. Using this observation and the σ -discreteness of the cover $\mathcal{U}_n$, show that $A_{n,m} = \bigcup_{U \in \mathcal{U}_n} U \times f_U^{-1}(W_m) \in \mathcal{A}_\alpha(X \times Y)$.

Now to get $f^{-1}(F) \in \mathcal{M}_{\alpha+1}(X \times Y)$ it suffices to verify that $f^{-1}(F) = \bigcap_{m=1}^\infty \bigcup_{n \geq m} A_{n,m}$.

To verify the inclusion $f^{-1}(F) \subset \bigcap_{m \in \mathbb{N}} \bigcup_{n \geq m} A_{n,m}$, fix any $m \in \mathbb{N}$ and $(x, y) \in X \times Y$ with $f(x, y) \in F$. Let $U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$, be a sequence of open sets containing the point x .

Since $s_n(U_n) \rightarrow x$ and f is continuous with respect to the first variable, we may find $k \geq m$ with $f(s_k(U_k), y) = f_{U_k}(y) \in W_m$. Thus $(x, y) \in U_k \times f_{U_k}^{-1}(W_m) \subset A_{k,m} \subset \bigcup_{n \geq m} A_{n,m}$ for every m .

Next, assume that $(x, y) \notin f^{-1}(F)$. Then $f(x, y) \notin F$ and there is $k \in \mathbb{N}$ with $\overline{W}_k \not\supset f(x, y)$. By the choice of the covers $\mathcal{U}_n$ and the continuity of f with respect to the first variable, there is $m \geq k$ such that $f(s_n(U_n), y) \notin \overline{W}_k$ for any $n \geq m$ and $x \in U_n \in \mathcal{U}_n$. Since $W_m \subset W_k$, we get $(x, y) \notin \bigcup_{n \geq m} A_{n,m}$. $\square$

4.2. Remark. The inclusion $CC(X \times Y, Z) \subset H_1(X \times Y, Z)$ is not true without restrictions on spaces X, Y, Z . If $X = Y = \{\infty\} \cup \Gamma$ is the one-point compactification of an uncountable discrete space Γ , then the function $f : X \times Y \rightarrow \mathbb{R}$ defined by

$$f(x, y) = \begin{cases} 1, & \text{if } x = y \in \Gamma; \\ 0, & \text{otherwise} \end{cases}$$

is separately continuous but does not belong to the class $H_1(X \times Y, \mathbb{R})$.

5. A GENERALIZATION OF THE RUDIN THEOREM

In [Ru] W. Rudin proved that every separately continuous function $f : X \times Y \rightarrow Z$ defined on the product of a metrizable space X and a topological space Y and acting into a locally convex topological vector space Z belongs to the first Baire class $B_1(X \times Y, Z)$. In [MMMS] it was proved that the metrizability of X in the Rudin Theorem can be replaced by the paracompactness and the σ -metrizability; moreover, if the space X is finite-dimensional then the local convexity of Z is superfluous, see [KM].

In this section we shall prove that the Rudin Theorem is still valid if X is replaced by any metrically quarter-stratifiable space and Z by a locally convex equiconnected space Z .

We remind that an *equiconnected space* is a pair (Z, λ) consisting of a topological space Z and a continuous map $\lambda : Z \times Z \times [0, 1] \rightarrow Z$ such that $\lambda(x, y, 0) = x$, $\lambda(x, y, 1) = y$, and $\lambda(x, x, t) = x$ for every $x, y \in Z$, $t \in [0, 1]$. For a subset $A \subset Z$ of an equiconnected space (Z, λ) let $\lambda^0(A) = A$ and $\lambda^n(A) = \lambda(\lambda^{n-1}(A) \times A \times [0, 1])$ for $n \geq 1$. Let also $\lambda^\infty(A) = \bigcup_{n \in \mathbb{N}} \lambda^n(A)$. An equiconnected space (Z, λ) is called *locally convex* if for every point $z \in Z$ and a neighborhood $O(z) \subset Z$ of z there is a neighborhood $U \subset Z$ of z with $\lambda^\infty(U) \subset O(z)$.

Equiconnected spaces are tightly connected with absolute extensors. We remind that a space Z is an *absolute extensor for a class $\mathcal{C}$* of topological spaces if every continuous map $f : B \rightarrow Z$ from a closed subspace B of a space $C \in \mathcal{C}$ admits a continuous extension $\bar{f} : C \rightarrow Z$ over all C . It is known that each equiconnected space is an absolute extensor for strongly countable-dimensional stratifiable spaces, while each locally convex equiconnected space is an absolute extensor for the class of stratifiable spaces. Moreover, a stratifiable space Z is an absolute extensor for stratifiable spaces if and only if Z admits an equiconnecting function λ turning Z into a locally convex equiconnected space, see [Bo], [Ca], [BGS].

Obvious examples of (locally convex) equiconnected spaces are convex subsets of (locally convex) linear topological spaces and their retracts. Also every contractible topological group G is an equiconnected space: an equiconnecting function λ on G can be defined by $\lambda(x, y, t) = h_t(xy^{-1}) \cdot h_t(e)^{-1} \cdot y$, where e is the neutral element of G and $\{h_t : G \rightarrow G\}_{t \in [0, 1]}$ is a contraction of G with $h_0 = \text{id}$ and $h_1(G) = \{e\}$. Let us recall that a topological space X is

contractible if it admits a homotopy $h : X \times [0, 1] \rightarrow X$ such that $h(x, 0) = x$ and $h(x, 1) = *$ for all $x \in X$ and some fixed point $*$ in X .

Given topological spaces X and Y denote by $C_p(X, Y)$ the subspace of the Tychonov product Y^X , consisting of all continuous functions from X to Y . Let $B_0(X, Y) = C_p(X, Y)$ and by transfinite induction for every ordinal $\alpha > 0$ define the Baire class $B_\alpha(X, Y)$ to be the sequential closure of the set $\bigcup_{\xi < \alpha} B_\xi(X, Y)$ in Y^X .

Now let us look at the Rudin theorem from the following point of view. Actually, this theorem states that every continuous function $f : X \rightarrow C_p(Y, Z)$ from a metrizable space X is a pointwise limit of “jointly continuous” functions. This observation leads to the following question: *Under which conditions every continuous map $f : X \rightarrow Z$ is a pointwise limit of some “nice” functions, and what should be understood under a “nice” function?*

In case of an equiconnected space (Z, λ) under “nice” functions we shall understand so-called piece-linear functions which are defined as follows. Let $PL(X, Z)$ be the smallest non-empty subset of $C_p(X, Z)$ satisfying the conditions:

- (1) for every point $z_0 \in Z$ and functions $f \in PL(X, Z)$, $\alpha \in C(X, [0, 1])$ the function $g(x) = \lambda(f(x), z_0, \alpha(x))$ belongs to $PL(X, Z)$;
- (2) a function $f : X \rightarrow Z$ belongs to $PL(X, Z)$ if there is an open cover $\mathcal{U}$ of X such that for every $U \in \mathcal{U}$ there is a function $g \in PL(X, Z)$ with $f|U = g|U$.

Equivalently, the set $PL(X, Z)$ can be defined constructively as the set of functions $f : X \rightarrow Z$ for which there is an open cover $\mathcal{U}$ of X such that for every $U \in \mathcal{U}$ there are points $z_0, \dots, z_n \in Z$ and functions $\alpha_1, \dots, \alpha_n \in C(X, [0, 1])$ such that $f|U = f_n|U$, where $f_0 \equiv z_0$ and $f_{i+1}(x) = \lambda(f_i(x), z_i, \alpha_i(x))$ for $0 \leq i < n$.

By $PL_{(1)}(X, Z)$ we denote the sequential closure of $PL(X, Z)$ in Z^X .

We recall that a topological space X is defined to be *strongly countable-dimensional* if X can be represented as the countable union $\bigcup_{n=1}^{\infty} X_n$ of closed subspaces with $\dim X_n < \infty$ for all n . According to [En₂, 5.1.10] a paracompact space X is strongly countable-dimensional if and only if X can be represented as the countable union $X = \bigcup_{n=1}^{\infty} X_n$ of closed subspaces such that every open cover $\mathcal{U}$ of X has an open refinement $\mathcal{V}$ with the property $\text{ord } \mathcal{V}|X_n \leq n$ for every $n \in \mathbb{N}$ (that is, every point $x \in X_n$ belongs to at most n elements of the cover $\mathcal{V}$).

5.1. Theorem. *Let X be a metrically quarter-stratifiable T_1 -space and (Z, λ) be an equiconnected space. If X is paracompact and strongly countable-dimensional or Z is locally convex, then $C_p(X, Z) \subset PL_{(1)}(X, Z)$.*

Proof. Fix any function $f \in C_p(X, Z)$. Using Theorem 2.2, find a weaker paracompact topology τ_p on X , a sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of τ_p -open covers of X and a sequence $\{s_n : \mathcal{U}_n \rightarrow X\}_{n \in \mathbb{N}}$ of functions such that $(x \in U_n \in \mathcal{U}_n, n \in \mathbb{N}) \Rightarrow (s_n(U_n) \rightarrow x)$. Since the space (X, τ_p) is paracompact, we may assume that each cover $\mathcal{U}_n$ is locally finite and contains no empty set.

If the space X is paracompact and strongly countable-dimensional, then according to the above-mentioned Theorem 5.1.10 of [En₂], we may additionally assume that X is represented as the union $X = \bigcup_{m=1}^{\infty} X_m$ of closed subsets such that $\text{ord } \mathcal{V}_n|X_m \leq m$ for every $n, m \in \mathbb{N}$.

For every $n \in \mathbb{N}$ take any partition of unity $\{\alpha_{n,U} : X \rightarrow [0, 1]\}_{U \in \mathcal{U}_n}$ subordinated to $\mathcal{U}_n$, that is $\alpha_{n,U}^{-1}(0, 1] \subset U$ for each $U \in \mathcal{U}_n$ and $\sum_{U \in \mathcal{U}_n} \alpha_{n,U} \equiv 1$.

Let $\xi_n : X \rightarrow N(\mathcal{U}_n)$, $\xi_n : x \mapsto \sum_{U \in \mathcal{U}_n} \alpha_{n,U}(x) \cdot U$, be the canonical map into the nerve of the cover $\mathcal{U}_n$, see [FF, VI.§3]. Next, we construct a map $\eta_n : N(\mathcal{U}_n) \rightarrow Z$ as follows. Denote

by $N^{(k)}(\mathcal{U}_n)$ the k -skeleton of $N(\mathcal{U}_n)$, where $k \geq 0$. The map η_n will be defined by induction. Let $\eta_n(U) = f(s_n(U))$ for every $U \in N^{(0)}(\mathcal{U}_n) = \mathcal{U}_n$. Suppose that η_n is already defined on the k -skeleton $N^{(k)}(\mathcal{U}_n)$ of $N(\mathcal{U}_n)$. We shall extend η_n onto $N^{(k+1)}(\mathcal{U}_n)$. Take any point $x \in N^{(k+1)}(\mathcal{U}_n) \setminus N^{(k)}(\mathcal{U}_n)$ and find a k -dimensional simplex $\sigma \ni x$. Fix any vertex v of σ . The point x can be uniquely written as $x = tv + (1-t)y$, where $t \in [0, 1]$ and $y \in \sigma \cap N^{(k)}(\mathcal{U}_n)$. Let $\eta_n(x) = \lambda(\eta_n(y), \eta_n(v), t)$ and observe that the so-extended map $\eta_n : N^{(k+1)}(\mathcal{U}_n) \rightarrow Z$ is continuous.

The inductive construction yields a continuous map $\eta_n : N(\mathcal{U}_n) \rightarrow Z$ which has the following property: $\eta_n(\sigma) \subset \lambda^\infty(\eta_n(\sigma^{(0)})) = \lambda^\infty(\bigcup_{U \in \sigma^{(0)}} f(s_n(U)))$ for every simplex σ of $N(\mathcal{U}_n)$. Observe also that $\eta_n \circ \xi_n \in PL(X, Z)$.

We claim that $\eta_n \circ \xi_n \rightarrow f$, provided X is strongly countable-dimensional or Z is locally convex. To show this, fix any point $x \in X$ and a neighborhood $O(f(x)) \subset Z$ of $f(x)$.

First, we consider the case when the space X is paracompact and strongly countable-dimensional. In this case $x \in X_m$ for some $m \in \mathbb{N}$ and $\text{ord } \mathcal{U}_n|_{X_m} \leq m$ for each n , which implies $\xi_n(x) \in N^{(m)}(\mathcal{U}_n)$ for every n . Using the continuity of the map λ , find a neighborhood $O_1 \subset Z$ of $f(x)$ such that $\lambda^m(O_1) \subset O(f(x))$. By the choice of the sequences $\{\mathcal{U}_n\}$ and $\{s_n\}$ there is $n_0 \in \mathbb{N}$ such that $s_n(U_n) \in f^{-1}(O_1)$ for every $n \geq n_0$ and $U_n \in \mathcal{U}_n$ with $U_n \ni x$. It follows from the construction of the map η_n that $\eta_n \circ \xi_n(x) \in \lambda^m(O_1) \subset O(f(x))$ for every $n \geq n_0$. Thus $\eta_n \circ \xi_n \rightarrow f$.

Now suppose that the equiconnected space (Z, λ) is locally convex. Then we may find a neighborhood $O_1 \subset Z$ of $f(x)$ with $\lambda^\infty(O_1) \subset O(f(x))$ and a number $n_0 \in \mathbb{N}$ such that $s_n(U_n) \in f^{-1}(O_1)$ for every $n \geq n_0$ and $x \in U_n \in \mathcal{U}_n$. It follows from the construction of the map η_n that $\eta_n \circ \xi_n(x) \in \lambda^\infty(O_1) \subset O(f(x))$ for every $n \geq n_0$. Thus in both cases the function sequence $\{\eta_n \circ \xi_n\}_{n=1}^\infty \subset PL(X, Z)$ tends to f , which proves that $f \in PL_{(1)}(X, Z)$. $\square$

Now we use Theorem 5.1 to generalize the Rudin Theorem as well as the results of [MMMS], [MS], [KMM], [KM]. By $CC(X \times Y, Z)$ the set of separately continuous functions $X \times Y \rightarrow Z$ is denoted.

5.2. Corollary. *Let X be a metrically quarter-stratifiable space, Y be a topological space, and (Z, λ) be an equiconnected space. If X is paracompact and strongly countable-dimensional or Z is locally convex, then $CC(X \times Y, Z) \subset B_1(X \times Y, Z)$.*

Proof. In obvious way the equiconnecting function λ induces an equiconnecting function on $C_p(Y, Z)$. Moreover, the equiconnected space $C_p(Y, Z)$ is locally convex if so is Z . Hence, we may apply Theorem 5.1 to conclude that $C_p(X, C_p(Y, Z)) \subset PL_{(1)}(X, C_p(Y, Z))$. Observe that the space $C_p(X, C_p(Y, Z))$ may be identified with $CC(X \times Y, Z)$, while every element of $PL(X, C_p(Y, Z))$ is jointly continuous as a function $X \times Y \rightarrow Z$. This yields that $CC(X \times Y, Z) \subset B_1(X \times Y, Z)$. $\square$

For an ordinal α and topological spaces X, Y, Z denote by $CB_\alpha(X \times Y, Z)$ the set of functions $f : X \times Y \rightarrow Z$ such that for every $x_0 \in X$ and $y_0 \in Y$ we have $f(\cdot, y_0) \in C(X, Z)$ and $f(x_0, \cdot) \in B_\alpha(Y, Z)$. Thus $CB_0(X \times Y, Z) = CC(X \times Y, Z)$. Generalizing the Rudin Theorem, V.K.Maslyuchenko et al. [MMMS] proved that $CB_\alpha(X \times Y, Z) \subset B_{\alpha+1}(X \times Y, Z)$ for any countable ordinal α , any metrizable space X , any topological space Y and any locally convex space Z , see [MMMS]. Below we have a further generalization.

5.3. Theorem. *Let X be a metrically quarter-stratifiable space, Y be a topological space and Z be a contractible space. Then $CB_\alpha(X \times Y, Z) \subset B_{\alpha+1}(X \times Y, Z)$ for every countable ordinal $\alpha > 0$.*

Proof. Let $\alpha \geq 1$ be a countable ordinal and $f \in CB_\alpha(X \times Y, Z)$. Fix a non-decreasing sequence (α_n) of ordinals with $\alpha = \sup_n \alpha_n + 1$ and let $h : Z \times [0, 1] \rightarrow Z$ be a contraction of Z with $h(z, 0) = z$ and $h(z, 1) = *$ for all $z \in Z$ and some fixed point $*$ in Z .

Using Theorem 2.2, find a weaker metrizable topology τ_m on X , a sequence $\{\mathcal{U}_n\}_{n \in \mathbb{N}}$ of τ_m -open cover of X and a sequence $\{s_n : \mathcal{U}_n \rightarrow X\}_{n \in \mathbb{N}}$ of maps such that $s_n(U_n) \rightarrow x$ for every $x \in U_n \in \mathcal{U}_n$, $n \in \mathbb{N}$. Using the paracompactness of (X, τ_m) , for every $n \in \mathbb{N}$ find a partition of unity $\{\alpha_{n,U} : (X, \tau_m) \rightarrow [0, 1]\}_{U \in \mathcal{U}_n}$ subordinated to $\mathcal{U}_n$, that is, $\alpha_{n,U}^{-1}(0, 1] \subset U$ for $U \in \mathcal{U}_n$. Then $\{\alpha_{n,U}^{-1}(0, 1]\}_{U \in \mathcal{U}_n}$ is a locally finite cover of X by functionally open sets.

Fix any well-ordering $\leq$ on the set $\mathcal{U}_n$. For every $U \in \mathcal{U}_n$ and $m \in \mathbb{N}$ consider the closed set $F_{m,U} = \alpha_{n,U}^{-1}[1/m, 1] \setminus \bigcup_{V < U} \alpha_{n,V}^{-1}(0, 1]$. Clearly, $\{F_{m,U}\}_{U \in \mathcal{U}_n}$ is a discrete collection of closed subsets of (X, τ_m) for every $m \in \mathbb{N}$. Using the collective normality of (X, τ_m) , for every $(m, U) \in \mathbb{N} \times \mathcal{U}_n$ we may find a continuous function $\beta_{m,U} : (X, \tau_m) \rightarrow [0, 1]$ such that $\beta_{m,U}^{-1}(1) \supset F_{m,U}$ and $\{\beta_{m,U}^{-1}(0, 1]\}_{U \in \mathcal{U}_n}$ is a discrete collection in (X, τ_m) for every $m \in \mathbb{N}$.

For every $U \in \mathcal{U}_n$ find a function sequence $\{f_{m,U}\}_{m \in \mathbb{N}} \subset \bigcup_{\xi < \alpha} B_\xi(Y, Z)$ with $\lim_{m \rightarrow \infty} f_{m,U}(y) = f(s_n(U), y)$ for every $y \in Y$. Without loss of generality, we may assume that $f_m \in B_{\alpha_m}(Y, Z)$ for every m . Then the function $g_{m,n} : X \times Y \rightarrow Z$ defined by the formula

$$g_{m,n}(x, y) = \begin{cases} h(f_{m,U}(y), 1 - \beta_{m,U}(x)), & \text{if } \beta_{m,U}(x) > 0 \text{ for some } U \in \mathcal{U}_n; \\ *, & \text{otherwise} \end{cases}$$

belongs to the class $B_{\alpha_m}(X \times Y, Z)$ for every $m, n \in \mathbb{N}$. Let us show that the limit $\lim_{m \rightarrow \infty} g_{m,n}(x, y)$ exists for every $(x, y) \in X \times Y$ and $n \in \mathbb{N}$. Indeed, given $(x, y) \in (X, Y)$ and $n \in \mathbb{N}$ let $U_n = \min\{U \in \mathcal{U}_n : \alpha_{n,U}(x) > 0\}$. Then $x \in F_{m,U_n}$ for all sufficiently large m and $g_{m,n}(x) = h(f_{m,U_n}(y), 1 - \beta_{m,U_n}(x)) = f_{m,U_n}(y) \rightarrow f(s_n(U_n), y)$ as $m \rightarrow \infty$. Thus $g_n(x, y) = \lim_{m \rightarrow \infty} g_{m,n}(x, y) = f(s_n(U_n), y)$ exists and $g_n \in B_\alpha(X \times Y, Z)$. Next, since $s_n(U_n) \rightarrow x$, we get $f(s_n(U_n), y) \rightarrow f(x, y)$ which yields that $f = \lim_{n \rightarrow \infty} g_n$ and $f \in B_{\alpha+1}(X \times Y, Z)$. $\square$

The following corollary generalizes a result from [MMMS].

5.4. Corollary. *Let $X_1, \dots, X_n$ be metrically quarter-stratifiable spaces, Y be a topological space, and (Z, λ) be an equiconnected space. If X_1 is paracompact and strongly countable-dimensional or Z is locally convex, then every separately continuous function $f : X_1 \times \dots \times X_n \times Y \rightarrow Z$ belongs to the n -th Baire class.*

We do not know if the conditions on X_1 or Z are essential.

5.5. Question. *Does every separately continuous function $f : X \times Y \rightarrow Z$ defined on the product of metrizable compacta and acting into a linear metric space belong to the first Baire class?*

5.6. Remark. Actually, according to [MMMS], the Rudin Theorem holds in a more strong form: $\bar{C}C(X \times Y, Z) \subset B_1(X \times Y, Z)$ for any metrizable space X , topological space Y , and a

locally convex space Z , where $\bar{C}C(X \times Y, Z)$ stands for the set of functions for which there is a dense subset $D \subset X$ such that for every $x_0 \in D$ and every $y_0 \in Y$ the functions $f(x_0, \cdot)$ and $f(\cdot, y_0)$ are continuous. It is interesting to remark that the metrizability of X is essential in this stronger form of the Rudin Theorem and can not be replaced by stratifiability; namely $\bar{C}C(A \times [0, 1], \mathbb{R}) \not\subset B_1(A \times [0, 1], \mathbb{R})$, where A is the Arens fan.

We recall the definition of the *Arens fan* A , see [En₁, 1.6.19]. Consider the following points of the real line: $a_0 = 0$, $a_n = \frac{1}{n}$ and $a_{nm} = \frac{1}{n} + \frac{1}{m}$, where $1 \leq n^2 \leq m$. Let $A_k = \{a_n : n \geq 0\} \cup \{a_{nm} : k \leq n \leq n^2 \leq m\}$ for $k \in \mathbb{N}$. On the union $A = \bigcup_{k \in \mathbb{N}} A_k$ consider the strongest topology inducing the original topology on each compactum A_k . Clearly, the so-defined space A is countable and stratifiable. The space A is known as an example of a sequential space which is not a Fréchet-Urysohn space.

5.7. Example. $\bar{C}C(A \times [0, 1], \mathbb{R}) \not\subset B_1(A \times [0, 1], \mathbb{R})$ for the Arens fan A .

Proof. We shall construct a function $f \in \bar{C}C(A \times [0, 1], \mathbb{R})$ which is not of the first Baire class. Take any function $f_0 \in B_2([0, 1], \mathbb{R}) \setminus B_1([0, 1], \mathbb{R})$. Write $f_0 = \lim_{n \rightarrow \infty} f_n$ where $\{f_n\}_{n \in \mathbb{N}} \subset B_1([0, 1], \mathbb{R})$. In its turn, represent each f_n as a pointwise limit $f_n = \lim_{m \rightarrow \infty} f_{nm}$ of continuous functions. Now consider the map $f : A \times [0, 1] \rightarrow \mathbb{R}$:

$$f(a, y) = \begin{cases} f_0(y), & \text{if } a = a_0; \\ f_n(y), & \text{if } a = a_n; \\ f_{nm}(y), & \text{if } a = a_{nm}. \end{cases}$$

It is easy to see that $f \in \bar{C}C(A \times [0, 1], \mathbb{R})$ but $f \notin B_1(A \times [0, 1], \mathbb{R})$. $\square$

6. RUDIN SPACES

In light of the mentioned generalizations of the Rudin Theorem it is natural to introduce the following

6.1. Definition. A topological space X is defined to be *Rudin* if for arbitrary topological space Y every separately continuous function $f : X \times Y \rightarrow \mathbb{R}$ belongs to the first Baire class.

In the following theorem we collect all fact concerning Rudin spaces we know at the moment. Let us recall that a subspace Y of a topological space X is *t-embedded*, if there is a continuous extender $E : C_p(Y) \rightarrow C_p(X)$, that is a map E such that $E(f)|_Y = f$ for every $f \in C_p(Y)$, see [Ar₁, III.§2.1]. Here $C_p(X) = C_p(X, \mathbb{R})$.

6.2. Theorem.

- (1) A space X is Rudin if and only if the calculation map $c_X : X \times C_p(X) \rightarrow \mathbb{R}$, $c_X : (x, f) \mapsto f(x)$, is of the first Baire class.
- (2) A space X is Rudin if the space $C_p(X)$ is Rudin.
- (3) Every metrically quarter-stratifiable space is Rudin.
- (4) If a space X is Rudin, then each *t-embedded* subspace of X is Rudin.
- (5) A space X is Rudin if $X = A \cup B$, where A, B are Rudin closed subspaces of X and A is a functionally closed retract of X .
- (6) If a Tychonov space X is Rudin, then $d(X) \leq l(X)$. Consequently, each compact Rudin space is separable.

Proof. 1) The “only if” part of the first statement is trivial. To prove the “if” part, assume that the calculation map $c : X \times C_p(X) \rightarrow \mathbb{R}$ is of the first Baire class. Observe that every separately continuous map $f : X \times Y \rightarrow \mathbb{R}$ can be seen as a continuous map $F : Y \rightarrow C_p(X)$ such that $f(x, y) = c(x, F(y))$ for $(x, y) \in X \times Y$. Now it is clear that f is of the first Baire class.

2) The second statement follows immediately from the first one.

3) According to Corollary 4.2, each metrically quarter-stratifiable space is Rudin.

4) Suppose X is a Rudin space and Y a t -embedded subspace of X . Let $E : C_p(Y) \rightarrow C_p(X)$ be the corresponding extender. Let $c_X : X \times C_p(X) \rightarrow \mathbb{R}$, $c_Y : Y \times C_p(Y) \rightarrow \mathbb{R}$ be the calculation functions for spaces X and Y , respectively. Taking into account that the function c_X is of the first Baire class and $c_Y(y, f) = c_X(y, E(f))$ for every $(y, f) \in Y \times C_p(Y)$ we conclude that c_Y is a function of the first Baire class either.

5) Suppose $X = A \cup B$, where A, B are Rudin closed subspaces of X and A is a functionally closed retract in X . Let $r : X \rightarrow A$ be a retraction and $\psi : X \rightarrow [0, 1]$ a map with $\psi^{-1}(0) = A$. Since the spaces A, B are Rudin, there are sequences of maps $\alpha_n : A \times C_p(A) \rightarrow \mathbb{R}$ and $\beta_n : B \times C_p(B) \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, tending to the calculation functions c_A and c_B of the spaces A, B , respectively. For every $n \in \mathbb{N}$ consider the function $\psi_n = \min\{n\psi, 1\}$ and define the map $c_n : X \times C_p(X) \rightarrow \mathbb{R}$ by the formula

$$c_n(x, f) = \begin{cases} \alpha_n(x, f|A), & \text{if } x \in A; \\ (1 - \psi_n(x))\alpha_n(r(x), f|A) + \psi_n(x)\beta_n(x, f|B), & \text{if } x \in B. \end{cases}$$

It is easy to see that the maps c_n are well-defined and continuous, and the sequence $\{c_n\}$ tends to the calculation function of X .

6) Suppose X is a Rudin space. Let $\{c_n\}_{n \in \mathbb{N}} \subset C(X \times C_p(X), \mathbb{R})$ be a sequence of continuous functions tending to the calculation function $c : X \times C_p(X) \rightarrow \mathbb{R}$. Then $c^{-1}(0) = \bigcap_{n \in \mathbb{N}} \bigcup_{m \geq n} c_m^{-1}(-1/n, 1/m)$ is a G_δ -set in $X \times C_p(X)$. Let $\{O_n\}_{n \in \mathbb{N}}$ be a sequence of open sets in $X \times C_p(X)$ with $\bigcap_{n=1}^\infty O_n = c^{-1}(0)$. By the continuity of the functions c_n , for every $x \in X$ and $n \in \mathbb{N}$ we may find a neighborhood $U_n(x)$ and a finite subset $F_n(x) \subset X$ such that $U_n(x) \times F_n(x)^\perp \subset O_n$, where $A^\perp = \{f \in C_p(X) : f|A \equiv 0\}$ for a subset $A \subset X$. For every $n \in \mathbb{N}$ choose a subset $X_n \subset X$ of size $|X_n| \leq l(X)$ with $\bigcup_{x \in X_n} U_n(x) = X$. We claim that $D = \bigcup \{F_n(x) : x \in X_n, n \in \mathbb{N}\}$ is a dense set in X . Otherwise, we would find a function $f \in C_p(X)$ with $f|D \equiv 0$ and $f(x) \neq 0$ for some $x \in X$. It is easy to see that $(x, f) \in \bigcap_{n \in \mathbb{N}} O_n$ but $(x, f) \notin c^{-1}(0)$, a contradiction, which shows that D is dense in X and $d(X) \leq |D| \leq l(X)$. $\square$

There are many open question about the structure of Rudin spaces.

6.3. Question. Is there a Rudin space which is not quarter-stratifiable? Is a (compact) space X Rudin, if $\{(x, f) : f(x) = 0\}$ is a G_δ -set in $X \times C_p(X)$? Is every (fragmentable, scattered or Rosenthal) compact Rudin space metrizable?

Let us remark that according to [Ar₂, II.6.1] each Corson Rudin compactum, being separable, is metrizable. Concerning the last question let us remark that there are non-metrizable compact spaces X such that $CC(X \times Y, \mathbb{R}) \subset B_1(X \times Y, \mathbb{R})$ for every compact space Y . The following result belongs to G.Vera [Ve].

6.4. Theorem. *A compact space X has countable Souslin number if and only if for every compact space Y every separately continuous function $f : X \times Y \rightarrow \mathbb{R}$ is of the first Baire class.*

Proof. Suppose X has a countable Souslin number and $f : X \times Y \rightarrow \mathbb{R}$ is a separately continuous function which can be seen as a continuous map $F : Y \rightarrow C_p(X)$. Then the image $F(Y) \subset C_p(X)$ is a compact subset of the space $C_p(X)$ over a compactum with countable Souslin number. According to Theorem 14 [Ar₁, II.§2] all such compact subsets are metrizable. By the Rudin Theorem, the restriction of the separately continuous calculation function $c : X \times C_p(X) \rightarrow \mathbb{R}$ onto $X \times F(Y)$ is of the first Baire class, which implies that the map $f(x, y) = c(x, F(y))$ is of the first Baire class either.

Now assume that the Souslin number of a compactum X is uncountable. Then there exists an uncountable family $\mathcal{U}$ of pairwise disjoint nonempty open sets in X . For every $U \in \mathcal{U}$ fix a continuous function $f_U : X \rightarrow [0, 1]$ such that $f_U|_{X \setminus U} \equiv 0$ and $\max f_U = 1$. Denote by 0 the origin of the vector space $C_p(X)$. It is easy to see that the set $Y = \{0\} \cup \{f_U : U \in \mathcal{U}\}$ in $C_p(X)$ is compact and the calculation function $c : X \times Y \rightarrow \mathbb{R}$, $c : (x, f) \mapsto f(x)$, is separately continuous. Yet, since $c^{-1}(1/2, 1]$ is not an F_σ -set in $X \times Y$, the function c is not of the first Baire class. In fact, one may show that $c \notin \bigcup_{\alpha < \omega_1} B_\alpha(X \times Y, \mathbb{R})$, i.e., c is not Baire measurable. $\square$

6.5. Remark. It follows from Theorems 6.2 and 6.4 that any unseparable compact space X with countable Souslin number is not Rudin but for every compact space Y every separately continuous function $f : X \times Y \rightarrow \mathbb{R}$ belongs to the first Baire class.

Finally let us prove one more result, connected with Baire classification of separately continuous functions.

6.6. Theorem. *Let X, Y, Z be compact Hausdorff spaces. If X is separable and Y has countable Souslin number, then every separately continuous function $f : X \times Y \times Z \rightarrow \mathbb{R}$ belongs to the second Baire class.*

Proof. Fix any separately continuous function $f : X \times Y \times Z \rightarrow \mathbb{R}$, which can be seen as a continuous map $F : X \rightarrow CC(Y \times Z, \mathbb{R})$ into the space of separately continuous functions endowed with the product topology. According to a recent result of S.Gulko and G.Sokolov [GS] if the space Y has countable Souslin number, then every compactum in $CC(Y \times Z, \mathbb{R})$ is Corson. Now if X is separable and Y has countable Souslin number, then $F(X) \subset CC(Y \times Z, \mathbb{R})$ is a separable Corson compactum. Since separable Corson compacta are metrizable, see [Ar₂, II.6.1], we conclude that the compactum $F(X)$ is metrizable. By [MMMS] (see also Corollary 5.4), the calculation map $c : F(X) \times Y \times Z \rightarrow \mathbb{R}$, $c : (g, y, z) \mapsto g(y, z)$, being separately continuous, belongs to the second Baire class. Then the function f belongs to the second Baire class because $f(x, y, z) = c(F(x), y, z)$. $\square$

6.6. Remark. The class of Rudin spaces includes the class of Lebesgue spaces introduced by O.Sobchuk in [So].

REFERENCES

- [Ar₁] A.V.Arkhangelskiĭ, *Topological function spaces*, MGU Publ., Moscow, 1989. (in Russian)
- [Ar₂] A.V.Arkhangelskiĭ, *Spaces of maps and rings of continuous functions*, Itogi Nauki i Tekhniki. General Topolog - 3, vol. 51, VINITI, Moscow, 1989, pp. 81–171. (in Russian)
- [Bo] C.Borges, *On stratifiable spaces*, Pacific J. Math. **17** (1966), 1–16.
- [Ca] R. Cauty, *Convexité topologique et prolongement des fonctions continues*, Compositio Math. **27** (1973), 233–271.
- [En₁] R.Engelking, *General Topology*, Mir, Moscow, 1986. (in Russian)
- [En₂] R.Engelking, *Theory of dimensions, finite and infinite*, Heldermann Verlag, Lemgo, 1995.
- [FF] V.V.Fedorchuk, V.V.Filippov, *general Topology. Basic Constructions*, MGU Publ., Moscow, 1988. (in Russian)
- [Gr] G.Gruenhage, *Generalized metric spaces*, Handbook of Set-Theoretic Topology (K.Kunen and J.Vaughan, eds.), Elsevier Sci., 1984, pp. 423–501.
- [GS] S.Gulko, G.Sokolov, *Compact spaces of separately continuous functions in two variables*, Topology Appl. **107** (2000), 89–96.
- [BGS] Bao-Lin Guo, K.Sakai, *Spaces of measures on stratifiable spaces*, Kobe J. Math. **12** (1995), 161–173.
- [KM] A.K.Kalancha, V.K.Maslyuchenko, *Baire classification of vector-valued separately continuous functions on products with finite-dimensional factor*, Collection of Sci. works of Kamjanets-Podilsky Ped. Univ. Ser. Phys.-Mat. **4** (1998), 43–46. (in Ukrainian)
- [KMM] A.K.Kalancha, V.K.Maslyuchenko, V.V.Mykhaylyuk, *Applications of Dugundji Theorem to problems Baire classification of vector-valued maps*, preprint. (in Ukrainian)
- [Ku₁] K. Kuratowski, *Sur la théorie des fonctions dans les espaces métriques*, Fund. Math. **17** (1931), 275–282.
- [Ku₂] K. Kuratowski, *Quelques problèmes concernant les espaces métriques non-séparables*, Fund. Math. **25** (1935), 534–545.
- [MMMS] V.K. Maslyuchenko, O.V. Maslyuchenko, V.V. Mykhaylyuk, O.V. Sobchuk, *Paracompactness and separately continuous mappings*, preprint.
- [MS] V.K.Maslyuchenko, O.V.Sobchuk, *Baire classification and σ -metrizable spaces*, Matem. Studii **3** (1994), 95–102. (in Ukrainian)
- [Mo] D. Montgomery, *Non-separable metric spaces*, Fund. Math. **25** (1935), 527–533.
- [Ny] P.J.Nyikos, *The theory of non-metrizable manifolds*, Handbook of Set-Theoretic Topology (K.Kunen and J.Vaughan, eds.), Elsevier Sci., 1984, pp. 633–684.
- [Ru] W.Rudin, *Lebesgue first theorem*, Math. Analysis and Applications, Part B. Adv. in Math. Supplem. Studies (Nachbin, ed.), vol. 78, Academic Press, 1981, pp. 741–747.
- [So] O. Sobchuk, *PP-spaces and Baire classification*, Intern. Conf. on Funct. Anal. and its Appl. (28–31 May, 2002), Lviv, p. 189.
- [St] A.H. Stone, *Paracompactness and product spaces*, Bull. Amer. Math. Soc. **54** (1948), 977–982.
- [Va] G.Vera, *Baire measurability of separately continuous functions*, Quart. J. Math. Oxford (2) **39** (1988), 109–116.